



# Cultural adaptation and Sepedi translation of the Activities-specific Balance Confidence scale



## Authors:

Tammy L. Prinsloo<sup>1</sup>   
Karin Joubert<sup>1</sup> 

## Affiliation:

<sup>1</sup>Department of Audiology,  
Faculty of Humanities,  
University of  
the Witwatersrand,  
Johannesburg, South Africa

## Corresponding author:

Karin Joubert,  
karin.joubert@wits.ac.za

## Dates:

Received: 19 July 2023

Accepted: 11 Feb. 2024

Published: 21 May 2024

## How to cite this article:

Prinsloo, T.L., & Joubert, K. (2024). Cultural adaptation and Sepedi translation of the Activities-specific Balance Confidence scale. *South African Journal of Communication Disorders*, 71(1), a1004. <https://doi.org/10.4102/sajcd.v71i1.1004>

## Copyright:

© 2024. The Authors.  
Licensee: AOSIS. This work  
is licensed under the  
Creative Commons  
Attribution License.

## Read online:



Scan this QR  
code with your  
smart phone or  
mobile device  
to read online.

**Background:** The Activities-specific Balance Confidence (ABC) scale is a widely used measure to identify older adults with balance difficulties. However, its applicability in the diverse South African context is hindered by cross-cultural and linguistic differences. Limited research exists on the use of the ABC scale in native South African languages.

**Objectives:** This study aimed to translate and culturally adapt the ABC scale into Sepedi, evaluate its reliability and determine self-perceived balance confidence among elderly individuals in a rural community.

**Method:** The ABC scale was translated and culturally adapted into Sepedi. Two trained raters administered the Sepedi version of the ABC (ABC-S) scale to 32 individuals aged between 60 and 88 years. Test-retest reliability and inter-rater reliability were determined, with one rater re-administering the scale 2 weeks later.

**Results:** Ten items from the original ABC scale were modified because of cultural, semantic or contextual inappropriateness. The ABC-S scale demonstrated very good intra- and inter-rater reproducibility, with an average intraclass correlation coefficient (ICC) of 0.85 and 0.81, respectively. The self-perceived balance confidence among elderly Sepedi individuals, as evaluated by the ABC-S scale, was high, with an average score of 81.3 and a range of 58.1 to 95.9.

**Conclusion:** The ABC-S scale is a reliable measurement tool to investigate balance confidence in Sepedi-speaking older adults.

**Contribution:** The ABC-S scale is a valuable screening tool for the identification of balance difficulties in Sepedi-speaking older adults as well as research settings.

**Keywords:** falls; aged; ABC scale; activities and participation; cultural adaptation; linguistic appropriateness.

## Introduction

Population ageing is a global phenomenon marked by an unprecedented growth of the elderly population (World Health Organization [WHO], 2022). The United Nations (2015a) observes that this population is growing more rapidly in developing countries compared to developed nations. Supporting this, the WHO (2022) predicts that four-fifths of older adults will be living in low- and middle-income countries in 2050, totalling approximately 1.2 billion elderly individuals (United Nations, 2019). In South Africa, the percentage of elderly people is on the rise, with the growth rate among those aged 60 years and older increasing from 1.1% in the year 2002–2003 to 3.0% in the year 2019–2020 (Statistics South Africa [Stats SA], 2020a). The rise is attributed to decreased fertility and increased longevity (Solanki et al., 2019), posing significant challenges to health and social systems, which are unprepared to address the chronic medical conditions associated with population ageing (WHO, 2022).

To enhance the quality of life of older adults and ensure that longer life is associated with positive experiences, the WHO (2002) suggests prioritising opportunities related to health, participation and security, a process known as 'active ageing' (p. 12). However, falls<sup>1</sup> and their related injuries often lead to fear of falling, loss of independence, institutionalisation (Khow & Visvanathan, 2017) and disability (Appeadu & Bordoni, 2023) and thus pose a threat to active ageing. Evidence suggests that near falls<sup>2</sup> are significant predictors of subsequent falls (Nagai et al., 2017), with falls accounting for the second most injury-related mortalities among the elderly globally (WHO, 2021).

1.A fall is defined as an event that results in a person inadvertently coming to rest on the ground or floor or other lower level (WHO, 2021).

2.A near fall, as defined by Maiden and colleagues (2014), is 'a stumble event or loss of balance that would result in a fall if sufficient recovery mechanisms were not activated' (p. 645).


An estimated 684 000 older adults worldwide suffer fatal falls annually (WHO, 2021). The global prevalence of falls among older adults is estimated to be 26.5% based on a systematic review of 104 articles (Salari et al., 2022). These statistics align with a study conducted in Cape Town, South Africa, where the prevalence rate of falls among the elderly was estimated to be 26.4% (Kalula et al., 2016). The United Nations (2015b) projects that the population of older adults in South Africa will double from 7.7 million in 2015 to 15.4 million in 2050, making falls a matter of concern.

Falls are complex events that result from the interactions between several risk factors (Chiarella et al., 2020). Elderly individuals face an increased risk of falling because of physical, sensory and cognitive changes associated with ageing (Montero-Odasso et al., 2022; WHO, 2021). Khow and Visvanathan (2017) categorise risk factors into two main groups: intrinsic and extrinsic factors. Intrinsic risk factors are specific to each individual and include functional abilities and comorbidities, such as the co-occurrence of hearing loss and vestibulopathy (Chiarella et al., 2020). Extrinsic risk factors, on the other hand, are external hazards such as dilapidated infrastructure contributing to the high prevalence rate of falls in South Africa (Kalula et al., 2016).

Despite national guidelines on fall prevention, there are no published data indicating their implementation in clinical practice (Kalu et al., 2019). In addition, there is a lack of preventative initiatives and educational programmes offered to the public in South Africa (Kalula et al., 2016), a concern given the potential role of various healthcare professionals in identifying individuals at risk of falling or with a fear of falling (De Clercq et al., 2020).

Audiologists, trained in the assessment and rehabilitation of vestibular problems, are well placed to assess fall risk, particularly because evidence suggests a link between hearing loss in older adults and increased fall risk (Jiam et al., 2016; Rogers, 2021). Audiologists, because of their access to older adults with hearing loss and vestibular problems, should thus play an integral role in screening of fall risk (Rogers, 2021), a critical step towards prevention of falls (Khow & Visvanathan, 2017). Despite this, the role of audiologists in fall risk assessment is not consistently emphasised across various countries' audiology scope of practice documents (Van Rie et al., 2022). There are a wide variety of screening measures to identify possible fall risk. A recent review conducted by Van Rie and colleagues (2022) reported that the four most reported fall risk screening measures used by audiologists, in order of frequency, were case history, the Timed Up and Go (TUG) test (Podsiadlo & Richardson, 1991), the Activities-specific Balance Confidence (ABC) scale (Powell & Myers, 1995) and the Dizziness Handicap Inventory (DHI) (Jacobson & Newman, 1990).

The 16-item ABC scale, developed by Powell and Myers (1995), is a reliable and valid (Botner et al., 2005; Huang & Wang, 2005; Salbach et al., 2006) measure widely used across



Source: World Health Organization. (2001). *International classification of functioning, disability and health*. Retrieved from <https://apps.who.int/iris/bitstream/handle/10665/42407/9241545429-eng.pdf?sequence=1&isAllowed=y>

FIGURE 1: International Classification of Functioning, Disability and Health.

multiple professions to identify older adults who may be at risk of falling by quantifying their self-perceived level of balance confidence in executing activities of daily living without falling or becoming unsteady (Sakakibara et al., 2011; Van Rie et al., 2022). In addition to use with community-dwelling elderly, the ABC scale is suitable for use with, among others, individuals with stroke (Botner et al., 2005; Seamon et al., 2019), vestibular disorders (Oh et al., 2023), persons with dystonia (Boyce et al., 2017) and Parkinson's disease (Franchignoni et al., 2014). This scale is quick to administer and correlates highly with the TUG, a functional mobility test, demonstrating the relationship between balance confidence and functional mobility (Hatch et al., 2003). The ABC scale further measures constructs within the International Classification of Functioning, Disability and Health (ICF) framework (Figure 1). The ICF is an internationally recognised framework that is underpinned by the biopsychosocial model of disability (Dantas et al., 2020). This comprehensive framework provides a scientific basis and common language for healthcare professionals worldwide to describe and classify falls, their associated risk factors in older adults (De Clercq et al., 2021), causes of falls and interventions aimed at fall prevention (Baris & Seren Intepeler, 2019).

In the activity domain, balance and falls gait (including stairs) and transfers are measured, while in the participation domain, it focuses on community function, home management, leisure and recreational activities role function and shopping (Academy of Neurologic Physical Therapy, 2013). This is important, as a recent study has found that fall assessment should focus on the activities and participation domains as this may decrease the risk of falling in older adults (De Clercq et al., 2021). These findings are supported by Johnson (2018) who explains that 'increased knowledge of the activities that are linked to falls and common fall-related injuries could be a valuable contribution to the prevention of falls in community-dwelling older adults' (p. 22).

The ABC scale has been adapted in other countries, such as Brazil (Marques et al., 2013), China (Hsu & Miller, 2006), Germany (Schott, 2008) and India (Moiz et al., 2017), among others. While the utility of the ABC scale has been recognised globally, it does not account for variations across different contexts, cultures or languages. South African citizens have the right to access healthcare services in a language of their

choice (Van den Berg, 2016). In the South African context, only one such adaptation (Kamanji, 2016) has occurred. Currently, no measures are available for assessing balance confidence in older Sepedi-speaking adults. When such measures are not available, health professionals may feel compelled to self-translate balance confidence scales, which may affect the reliability thereof (Sahu & Kulkarni, 2017).

In South Africa, Sepedi is the third largest African language, the fifth most spoken language in the country and the first language of approximately 4.6 million people (Stats SA, 2011). The majority (52.9%) of first language Sepedi speakers live in the Limpopo province. Sepedi is also spoken by 10.6% of people who live in Gauteng and 9.3% of people who live in Mpumalanga (Stats SA, 2011). Therefore, the aim of this study was to translate and culturally adapt the ABC scale into Sepedi, as well as determine the reliability of the Sepedi version of the ABC (ABC-S) scale. The secondary aim was to determine the self-perceived balance confidence of first language Sepedi speakers in a rural community in Limpopo.

## Methodology

### Research design

A mixed-methods research approach was employed for this two-phased study. In the first phase, a formative qualitative methodology was used to translate and culturally adapt the ABC scale (WHO, 2016). In the second phase, a descriptive research design was used to determine the reliability of the ABC-S scale and describe the self-perceived balance confidence among first language Sepedi speakers in a rural community in Limpopo.

### Context

The second phase of this study was conducted in a rural area of the Greater Sekhukhune District Municipality of the Limpopo province, where Sepedi is the predominant language spoken, accounting for 83% of the population (Integrated Development Plan, 2021).

### Participants

Using purposive sampling, a total of 32 participants were recruited for the second phase of the study. Participants were required to meet the following criteria: (1) be between the ages of 60 and 88 years (Kamanji, 2016; Marques et al., 2013) to allow for comparison with the isiZulu (Kamanji, 2016) and Brazilian (Marques et al., 2013) versions of the scale; (2) be first language Sepedi speakers; (3) reside in a rural community; (4) be ambulatory and capable of independently performing daily activities inside and outside of the home either with or without walking aids; and (5) be able to understand and follow multistep instructions. Individuals with medical conditions that increase the risk of falling (e.g. neurological disorders, musculoskeletal disorders, orthostatic hypotension, etc.) as well as those with unmanaged hearing and visual impairments were excluded from the study.

The mean age of participants was 70.06 years ( $\pm 7.26$ ; range: 60–88) and the majority were female ( $n = 84\%$ ). The gender distribution aligns with the distribution of the broader Sekhukhune District Municipality population where 65% of the population in this age group is female (Integrated Development Plan, 2021).

### Measures

Data collection involved the use of a self-developed case history questionnaire (Appendix 1) and the ABC-S scale. The case history questionnaire consisted of 19 questions organised in four sections: demographic information, balance history, medical history and family history. The development of the case history questionnaire was guided by literature (Kamanji, 2016). The ABC-S scale has 16 items rated on an 11-point subscale ranging from 0% (no confidence) to 100% (completely confident) with increments of 10%. Scores below 50% indicate a low level of physical functioning. Scores above 80% represent a high level of physical functioning (Marques et al., 2013). Scores of <67% have shown to be predictive of future falls (Reelick et al., 2009).

### Data collection

#### First phase

The first phase of data collection involved the cultural adaptation and translation of the ABC scale into Sepedi, following a five-step process as outlined by the WHO (2016). This process included forward translation, expert panel assessment, back translation, pretesting and final version development.

#### Forward translation

The ABC scale was translated into Sepedi by a translation agency with 26 years' experience in healthcare and medical translations. All translators hired by the agency are native speakers with the relevant credentials. The translator, appointed by the agency, was a first language Sepedi speaker proficient in English.

#### Expert panel assessment

The expert panel comprised three health professionals (two nurses and one pharmacist) with an undergraduate degree and a minimum of 3 years' experience in working with older adults in rural communities. They were purposively selected from a health facility in the area and resided in the community where the study was conducted. They were requested to rate the cultural and contextual relevance of each activity assessed in the scale to the rural South African context. Each question was rated for relevance, with scores ranging from 1 (highly irrelevant) to 5 (highly relevant). For questions judged to be culturally or contextually irrelevant (ratings of 1 and 2), panel members were required to provide an explanation and a more culturally or contextually relevant alternative. Items with a score of 4 (relevant) and 5 (highly relevant) were considered culturally and contextually relevant. An inter-judge percentage agreement of  $\geq 80\%$  was used (McHugh, 2012).

### Back translation

The culturally adapted ABC-S scale was translated back into English by an independent translator (proficient in Sepedi and English) from the same translation agency used in the forward translation.

### Pretesting

The ABC-S scale was piloted on three female participants, aged 65, 72 and 80 years, respectively, and who met the same inclusion criteria as for the main study. The objectives of the pilot study were to determine the semantic, contextual and cultural appropriateness of each question in the ABC-S scale. Eight modifications were made based on the feedback from the pilot study participants that led to the final version of the ABC-S scale (Appendix 2).

### Second phase

During the second phase, two research assistants were trained to recruit participants, complete the case history questionnaire and administer the ABC-S scale. The research assistants were employed as hearing screeners in the community where the study was conducted and had at least 4 years' experience in case history taking. They were both first language Sepedi speakers proficient in English. Potential participants were recruited at two of the local primary healthcare clinics using purposive sampling techniques. Only those who provided written informed consent were included in the study. Data collection occurred over an 8-week period, either at participants' homes or community sites based on their preferences. The ABC-S scale was administered by the two research assistants during the initial data collection period and readministered 2 weeks later by rater one to assess inter-rater and test-retest reliability.

Data were encoded and analysed using appropriate statistical techniques. The Cronbach's alpha coefficient and item-total correlations were utilised to assess the internal consistency of the ABC-S scale. Test-retest reliability and inter-rater reliability were determined using the intraclass correlation coefficient (ICC). Descriptive statistics, including mean, range and standard deviation, were used to describe the self-perceived balance confidence.

### Ethical considerations

Ethical clearance to conduct this study was obtained before data collection from the University of the Witwatersrand School of Human and Community Development Ethics Committee (No. STA\_2021\_5). The research adhered to the ethical principles outlined in the World Medical Association Declaration of Helsinki (World Medical Association [WMA], 2013). Permission to conduct the study was obtained from relevant authorities. Participants were fully informed of the nature of the study as well as their rights and assured of anonymity and confidentiality. Participants identified as having a fall risk were referred to their closest audiology department for assessment. Three options were available: the local community-based audiology clinic or two hospitals in

**TABLE 1:** Original and modified items in the cross-cultural adaptation of the Activities-specific Balance Confidence scale.

Item no.	Original items of the ABC scale	Modified items of the ABC-S scale
3	Bend over and pick up a slipper from the front of a closet floor	Bend over and pick up a shoe from the area where one stores their belongings
4	Reach for a small can off a shelf at eye level	Reach for a small tin that is on top of a shelf that you can see
8	Walk outside the house to a car parked in the driveway	Walk outside to a vehicle in the yard
9	Get into or out of a car	Get into or out of a vehicle
10	Walk across a parking lot to the mall	Walk across a parking lot to the mall or shops
11	Walk up or down a ramp	Walk up or down a hill
12	Walk in a crowded mall where people rapidly walk past you	Walk in a crowded mall or shop where people rapidly walk past you
13	Are bumped into by people as you walk through the mall	Are bumped into by people as you walk through the mall or shops
15	Step onto or off an escalator while holding onto parcels such that you cannot hold onto the railing	Step onto or off an escalator while holding onto luggage such that you cannot hold onto the railing
16	Walk outside on icy sidewalks	Walk outside on surfaces largely covered by dew

ABC, Activities-specific Balance Confidence; ABC-S, Sepedi version of the ABC.

the Greater Sekhukhune District Municipality. The two hospitals also offer physiotherapy services.

## Results and discussion

The aims of this study were to translate and culturally adapt the ABC scale into Sepedi as well as determine the reliability of the ABC-S scale. In addition, the self-perceived balance confidence of first language Sepedi speakers in a rural community in Limpopo was determined.

### Cross-cultural adaptation of the translated Activities-specific Balance Confidence scale

The translated ABC scale (Appendix 2) was culturally adapted by a bilingual expert panel to suit the needs of the rural South African context. Ten of the original 16 items were modified to ensure semantic, contextual and cultural appropriateness. These modifications are documented in Table 1.

The adjustments were necessary because of the distinct local reality and cultural differences of the South African participants, considering that the original scales development is based on the Canadian population (Powell & Myers, 1995).

For instance, item 3 originally 'pick up a slipper from the front of the closet floor' underwent modifications to 'pick up a shoe from the area where one stores their belongings' to better align with local terminology and cultural context. Specifically, the terms 'slipper' and 'closet' were not familiar to the South African participants. Marques et al. (2013) caution that using unfamiliar terminology may hinder comprehension. None of the participants reported any doubts as to the meaning during the administration of the ABC-S scale. The adaptations aimed to ensure comprehension without altering the activity's intended meaning, as evidenced by comparable scores to the Brazilian-Portuguese and Canadian studies (Table 5).

Similarly, item 4 was also adapted by changing the phrase 'reach for a small can off a shelf at eye level' to 'reach for a small tin that is on top of a shelf that you can see' to align with the common expression in the Sepedi culture.

Items 8 and 9 were adapted to account for socioeconomic differences, replacing 'walk outside the house to a car parked in the driveway' with 'walk outside to a vehicle in the yard' emphasising 'vehicle' to accommodate varied transportation modes. Most of the residents in Limpopo use taxis (Stats SA, 2020b), as cars are owned by less than 15% of households in rural communities (Stats SA, 2018). Driveways are also uncommon in rural communities.

To align with the contextual realities of the Elias Motsoaledi Local Municipality (EMLM) area and the lifestyle of its residents, the term 'shop(s)' was added to items 10, 12 and 13. Most of the people who reside in the EMLM area and other rural areas in South Africa go to shops, which are more accessible to them compared to malls.

During the pilot study, participants expressed unfamiliarity with the term 'ramp' in item 11 ('walk up or down a ramp'), leading to its modification 'walk up or down a hill'. In item 15, the word 'parcels' was replaced with 'luggage' to better suit the context of the study population.

Item 16 ('walk outside on icy sidewalks') underwent the final modification to account for differences in climate, as snow is extremely rare in South Africa. The term 'sidewalk' was also changed. Item 16 was thus changed to 'walk outside on a surface largely covered by dew' to reflect context and align with the Sepedi expression.

These meticulous adaptations underscore the significance of tailoring assessment tools for cultural relevance, ensuring the reliability of the ABC-S scale in the local South African setting.

### Reliability of the Sepedi version of the ABC scale

The test-retest reliability of the ABC-S scale is shown in Table 2. The ICC for the total score exhibited a high level of reliability, measuring at 0.85, with a 95% confidence interval. This is consistent with the results obtained for the isiZulu ABC scale, which demonstrated a score of 0.82 (Kamanji, 2016) and balance confidence in individuals with stroke at 0.85 (Botner et al., 2005). Other versions of the ABC scale have demonstrated excellent test-retest reliability, such as the Brazilian-Portuguese (Marques et al., 2013), Hindi (Moiz et al., 2017) and Chinese (Hsu & Miller, 2006) versions with scores of 0.94, 0.97 and 0.99, respectively.

Table 3 shows the inter-rater reliability of the ABC-S scale. The ICC for the total score exhibits a high level of reliability, measuring at 0.81 with a confidence interval of 95%. This is consistent with other versions of the ABC scale such as the

**TABLE 2:** Test-retest reproducibility: Mean (standard error) and intraclass correlation coefficient of the Sepedi version of the Activities-specific Balance Confidence scale.

ABC-S scale items	Evaluation 1		Evaluation 2	
	Mean <sup>†</sup>	SE	Mean <sup>‡</sup>	SE
1	80	20.0	97.5	2.5
2	67.5	6.3	52.5	22.1
3	65	15.6	60	19.6
4	70	10.0	72.5	24.3
5	57.5	13.8	62.5	22.5
6	30	17.8	20	9.1
7	85	2.9	90	4.1
8	92.5	4.8	97.5	2.5
9	87.5	7.5	100	0.0
10	75	16.6	77.5	16.5
11	75	10.4	57.5	19.3
12	55	11.9	80	16.8
13	67.5	16.5	80	16.8
14	55	20.2	22.5	22.5
15	45	18.5	20	20.0
16	100	0.0	95	5.0
<b>Total</b>	<b>1107.5</b>	<b>-</b>	<b>1085.0</b>	<b>-</b>

Note: Average ICC = 0.85.

ICC, intraclass correlation coefficient; ABC-S, Sepedi version of the Activities-specific Balance Confidence scale.

<sup>†</sup>, average = 69.2; <sup>‡</sup>, average = 67.8.

**TABLE 3:** Inter-rater reproducibility: Mean (standard error) and intraclass correlation coefficient of the Sepedi version of the Activities-specific Balance Confidence scale.

ABC-S scale items	Rater 1		Rater 2	
	Mean <sup>†</sup>	SE	Mean <sup>‡</sup>	SE
1	83.75	8.9	56.25	11.3
2	87.50	7.7	77.50	8.0
3	88.75	5.2	82.50	5.9
4	85.00	8.9	75.00	7.6
5	87.50	6.8	80.00	6.6
6	76.25	7.3	67.50	6.5
7	96.25	2.6	87.50	4.9
8	95.00	3.8	87.50	5.6
9	100.00	0.0	95.00	1.9
10	96.25	2.6	90.00	4.2
11	95.00	2.7	88.75	5.5
12	97.50	2.5	88.75	4.8
13	97.50	1.6	87.50	3.1
14	85.00	5.0	75.00	6.8
15	82.50	7.3	71.25	7.7
16	100.00	0.0	92.50	3.1
<b>Total</b>	<b>1453.75</b>	<b>-</b>	<b>1313.75</b>	<b>-</b>

Note: Average ICC = 0.81.

ICC, intraclass correlation coefficient; ABC-S, Sepedi version of the Activities-specific Balance Confidence scale.

<sup>†</sup>, average = 90.86; <sup>‡</sup>, average = 81.41.

Brazilian-Portuguese translation, with a score of 0.80 (Marques et al., 2013). These findings suggest that the scale produces consistent results over time and between different raters and can be used in clinical and research settings.

Table 4 shows the internal consistency of the ABC-S scale, which was determined using the Cronbach's alpha coefficient and item-total correlations. The scale showed a high internal consistency of  $\alpha = 0.92$ . This was reinforced by moderate-to-strong item-total correlations for 13 of the items, ranging from 0.41 to 0.85 and exceeding the cutoff value of 0.70 (Ferketich, 1991).

An  $\alpha \geq 0.9$  is considered excellent (Bland & Altman, 1997). This indicates that the items of the ABC-S scale measure a single, unidimensional construct (Tavakol & Dennick, 2011) and is thus appropriate for use in clinical and research settings. The internal consistency of the ABC-S scale is consistent with findings reported in previous studies (see Guan et al., 2012; Kamanji, 2016; Mak et al., 2007; Powell & Myers, 1995; Salbach et al., 2006; Schott, 2008; Van Heuvelen et al., 2005). Subsequent analysis indicated that the majority of the items are suitable for retention, leading to a decrease in the alpha coefficient if they are excluded. Nevertheless, there were four exceptions to this trend, specifically items 1, 3, 10 and 16, for which the removal would result in an increased alpha coefficient of 0.92. Therefore, it is advisable to consider removing these items. This aligns with the observed item-total correlations for items 1, 3 and 16, which demonstrated a

weak correlation. According to Powell and Myers (1995), the internal consistency of the scale does not exhibit a significant improvement when a few items are removed. Thus, the deletion of item 16, which pertains to 'walking outside on icy sidewalks', should not pose any challenges for administering the scale in warmer climates, such as South Africa (Myers et al., 1998).

### Self-perceived balance confidence of elderly people

The mean confidence level of first language Sepedi speakers in a rural community in the Limpopo province was 81.3 ( $\pm 10.3$ ; range: 58.1–95.9) (Table 5). This is suggestive of a high level of self-perceived confidence in their ability to maintain balance during various daily activities (Myers et al., 1998).

Subjective fear of falling was assessed through the unvalidated single-item question: 'Do you have a fear of falling?' asked during case history. Of the 12 participants who indicated fear of falling, 6 (50%) obtained a total score of <67, with scores ranging from 23.13 to 65. Only two participants (6.25%) who did not report a fear of falling obtained scores below 67. The single-item question was significantly correlated to the ABC scores in the sample ( $r = -0.567$ ).

The two items with the lowest degrees of confidence in this study were identical to the findings in the Brazilian (Marques et al., 2013), isiZulu (Kamanji, 2016) and Canadian populations (Powell & Myers, 1995). These items generally involve activities that pose greater challenges or perceived risks for the participants. These include item 6 ('stand on a chair and reach for something') and item 15 ('step onto or off an escalator while holding onto parcels such that you cannot hold onto the railing'). These items had values ranging from 58.1 to 64.4,

**TABLE 4:** Internal consistency of the Sepedi version of the Activities-specific Balance Confidence scale.

Item no.	Corrected item-total correlation	Alpha, if item deleted
1	0.26	0.92
2	0.68	0.91
3	0.29	0.92
4	0.66	0.91
5	0.74	0.91
6	0.76	0.91
7	0.65	0.91
8	0.64	0.91
9	0.65	0.91
10	0.41	0.92
11	0.85	0.90
12	0.68	0.91
13	0.62	0.91
14	0.77	0.90
15	0.78	0.90
16	0.30	0.92

**TABLE 5:** Comparison of the Sepedi version of the Activities-specific Balance Confidence score with the Brazilian, isiZulu and the original English version.

Item no.	ABC-S score (n = 32)		Brazilian ABC score (n = 40)		isiZulu ABC score (n = 32)		Original ABC score (n = 60)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
1	89.4	-	88.2	-	74.2	-	87.5	-
2	80.0	-	85.5	-	48.8	-	64.8	-
3	87.5	-	86.2	-	57.9	-	62.8	-
4	77.8	-	93.5	-	60.6	-	89.5	-
5	79.7	-	80.5	-	47.9	-	46.5	-
6	58.1	-	68.5	-	26.9	-	38.0	-
7	88.4	-	96.7	-	63.0	-	66.8	-
8	89.7	-	87.2	-	72.7	-	71.6	-
9	90.6	-	86.5	-	68.2	-	78.5	-
10	83.4	-	89.7	-	48.5	-	67.7	-
11	81.6	-	81.5	-	41.2	-	61.0	-
12	82.2	-	84.0	-	51.2	-	62.2	-
13	84.1	-	79.5	-	41.2	-	53.1	-
14	67.2	-	76.7	-	49.1	-	52.3	-
15	64.4	-	63.2	-	28.8	-	31.3	-
16	95.9	-	61.2	-	36.1	-	20.7	-
<b>Total score</b>	<b>81.3</b>	<b>10.3</b>	<b>81.7</b>	<b>10.1</b>	<b>51.0</b>	<b>14.4</b>	<b>59.6</b>	<b>27.7</b>

Source: Adapted from Kamanji, R. (2016). Zulu translation and cross-cultural adaptation of the activities-specific balance confidence (ABC) scale. Research report. University of the Witwatersrand; Marques, A.P., Mendes, Y.C., Taddei, U., Pereira, C.A., & Assumpção, A. (2013). Brazilian-Portuguese translation and cross-cultural adaptation of the activities-specific balance confidence (ABC) scale. *Brazilian Journal of Physical Therapy*, 17(2), 170–178. <https://doi.org/10.1590/S1413-35552012005000072>; Powell, L.E., & Myers, A.M. (1995). The activities-specific balance confidence (ABC) scale. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 50A(1), M28–M34. <https://doi.org/10.1093/gerona/50A.1.M28>

Note: Items were rated from 0% (no confidence) to 100% (complete confidence).

ABC-S, Sepedi version of the ABC; ABC, Activities-specific Balance Confidence.

consistent with findings from other studies (Botner et al., 2005; Mak et al., 2007; Salbach et al., 2006; Schepens et al., 2010). Item 16 in this study, however, had the highest score (95.9), in contrast to the Canadian and Brazilian population at 20.7 and 61.2, respectively. This difference may be explained by the perceived level of difficulty walking on a surface covered by dew (this study) in comparison to walking on ice.

A score lower than 67% has been identified as an indicator of fall risk (Lockhart et al., 2019; Reelick et al., 2009). When comparing the mean scores across different cultural and linguistic contexts, it is evident that the Sepedi (81.3) and Brazilian-Portuguese (81.7) studies yielded considerably higher mean scores compared to the isiZulu (51) and Canadian (59.6) studies, as shown in Table 5. In both the Sepedi and Brazilian-Portuguese studies, only three items obtained scores lower than 67%, indicating a relatively low fall risk among the participants. This finding is in stark contrast to the isiZulu (Kamanji, 2016) and Canadian (Powell & Myers, 1995) versions of the ABC scale, where 13 and 12 items, respectively, received scores below the fall risk threshold. These results suggest that elderly individuals in the Sepedi and Brazilian-Portuguese-speaking populations demonstrated a higher level of self-perceived balance confidence when compared to their Canadian counterparts, and a lower likelihood of falling when compared to their isiZulu and Canadian counterparts.

The disparity on fall risk among these populations may be attributed to differences in the selection criteria employed in the original and isiZulu studies. In a study conducted by Lajoie and Gallagher (2004), individuals who had experienced a stroke or had preexisting decreased postural control were included as participants. This selection criterion likely contributed to lower balance confidence scores in the Canadian study population. Similarly, the isiZulu study by Kamanji (2016) included participants with physical impairments. These impairments may have negatively impacted balance confidence levels, resulting in the comparatively lower mean scores observed in the isiZulu study. In contrast, the current and Brazilian-Portuguese studies (Marques et al., 2013) excluded individuals with medical conditions that increase the risk of falling and those with unmanaged hearing and visual impairments. Consequently, the participants in these studies may have had a higher baseline level of balance confidence, leading to the elevated mean scores observed.

## Conclusion

The adapted ABC-S scale demonstrated high internal consistency, test-retest reliability and inter-rater reliability. The findings of this study support the recommendation for the use of the ABC-S scale in Sepedi-speaking older adults to assess their self-perceived level of balance confidence. It is a valuable tool for clinicians, researchers and policymakers working in the field of fall prevention that can contribute to the development and implementation of effective preventative initiatives, ultimately improving the overall well-being and quality of life of Sepedi-speaking older adults. Furthermore, the cultural adaptation of the ABC

scale into Sepedi enhances its applicability and ensures that it resonates with the linguistic and cultural contexts of the target population. This consideration is crucial in promoting inclusivity and sensitivity to diverse cultural backgrounds (Pascoe & Norman, 2011) ultimately leading to more accurate and meaningful assessments of balance confidence.

A limitation of the study is the small sample size and the recruitment of participants from only one municipal area in the Limpopo province, which limits the generalisability of the study. Further studies should explore the validity of the tool as well as the replication of the study across the different provinces and contexts (e.g., urban vs. rural) where significant numbers of Sepedi-speaking individuals reside as well as medical conditions. This will assist with the generalisation of the results. The ABC scale should also be translated and culturally adapted into other indigenous South African languages.

## Acknowledgements

The authors would like to thank all the individuals who participated in the study. A special thank you to the research assistants for their significant contribution to the study.

## Competing interests

The authors have declared that no competing interest exists.

## Authors' contributions

T.L.P. and K.J. were involved in the conceptualisation of the project, methodology, writing and reviewing of the manuscript. T.L.P. was responsible for data collection and analysis.

## Funding information

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

## Data availability

The authors confirm that the data supporting the findings of this study are available within the article and its supplementary materials.

## Disclaimer

The views and opinions expressed in this article are those of the authors and do not necessarily reflect the official policy or position of any affiliated agency of the authors, or that of the publisher. The authors are responsible for this article's results, findings, and content.

## References

- Academy of Neurologic Physical Therapy. (2013). *Vestibular edge outcome measures taskforce*. Retrieved from <https://www.neuropt.org/docs/default-source/ vestibular-edge/ vestibular-edge-outcomes-measures.pdf?sfvrsn=a102>
- Appeadu, M.K., & Bordon, B. (2023). Falls and fall prevention in older adults. In *StatPearls*. StatPearls Publishing. Retrieved from <https://www.ncbi.nlm.nih.gov/ books/NBK560761/>

- Baris, V.K., & Seren Intepeler, S. (2019). Views of key stakeholders on the causes of patient falls and prevention interventions: A qualitative study using the international classification of functioning, disability and health. *Journal of Clinical Nursing*, 28(3–4), 615–628. <https://doi.org/10.1111/jocn.14656>
- Bland, J.M., & Altman, D.G. (1997). Statistics notes: Cronbach's alpha. *British Medical Journal*, 314(7080), 570–572. <https://doi.org/10.1136/bmj.314.7080.572>
- Botner, E.M., Miller, W.C., & Eng, J.J. (2005). Measurement properties of the activities-specific balance confidence scale amongst individuals with stroke. *Disability and Rehabilitation*, 27(4), 156–163. <https://doi.org/10.1080/09638280400008982>
- Boyce, M.J., Lam, L., Chang, F., Mahant, N., Fung, V.S.C., & Bradnam, L. (2017). Validation of fear of falling and balance confidence assessment scales in persons with dystonia. *Journal of Neurologic Physical Therapy*, 41(4), 239–244. <https://doi.org/10.1097/NPT.0000000000000198>
- Chiarella, G., Pisani, D., & Viola, P. (2020). Disequilibrium and risk of falling in the elderly is a priority for health services. *Reviews on Recent Clinical Trials*, 15(3), 162–163. <https://doi.org/10.2174/1574887115666200630105529>
- Dantas, D.D.S., Correa, A.P., Buchalla, C.M., Castro, S.S.D., & Castaneda, L. (2020). Biopsychosocial model in health care: Reflections in the production of functioning and disability data. *Fisioterapia em Movimento*, 33, e003321. <https://doi.org/10.1590/1980-5918.033.a021>
- De Clercq, H., Naude, A., & Bornman, J. (2020). The perspectives of healthcare practitioners on fall risk factors in older adults. *Health SA Gesondheid*, 25, a1495. <https://doi.org/10.4102/hsag.v25i0.1495>
- De Clercq, H., Naudé, A., & Bornman, J. (2021). Older adults' perspectives on fall risk: Linking results to the ICF. *Journal of Applied Gerontology*, 40(3), 328–338. <https://doi.org/10.1177/0733464820929863>
- Ferketich, S. (1991). Focus on psychometrics: Aspects of item analysis. *Research in Nursing & Health*, 14(2), 165–168. <https://doi.org/10.1002/nur.4770140211>
- Franchignoni, F., Giordano, A., Ronconi, G., Rabini, A., & Ferriero, G. (2014). Rasch validation of the activities-specific balance confidence scale and its short versions in patients with Parkinson's disease. *Journal of Rehabilitation Medicine*, 46(6), 532–539. <https://doi.org/10.2340/16501977-1808>
- Guan, Q., Han, H., Li, Y., Zhao, L., Jin, L., & Zhan, Q. (2012). Activities-specific balance confidence (ABC) scale adapted for the mainland population of China. *Clinical Rehabilitation*, 26(7), 648–655. <https://doi.org/10.1177/0269215511427748>
- Hatch, J., Gill-Body, K.M., & Portney, L.G. (2003). Determinants of balance confidence in community-dwelling elderly people. *Physical Therapy*, 83(12), 1072–1079. <https://doi.org/10.1093/ptj/83.12.1072>
- Hsu, P.C., & Miller, W.C. (2006). Reliability of the Chinese version of the activities-specific balance confidence scale. *Disability and Rehabilitation*, 28(20), 1287–1292. <https://doi.org/10.1080/09638280600638414>
- Huang, T.T., & Wang, W.S. (2009). Comparison of three established measures of fear of falling in community-dwelling older adults: Psychometric testing. *International Journal of Nursing Studies*, 46(10), 1313–1319. <https://doi.org/10.1016/j.ijnurstu.2009.03.010>
- Integrated Development Plan. (2021). *Draft IDP 2021/2022: Sekhukhune District Municipality*. Retrieved from <https://www.sekhukhunedistrict.gov.za/sdm-admin/documents/DRAFT%20IDP%20202122.pdf>
- Jacobson, G.P., & Newman, C. (1990). The development of the dizziness handicap inventory. *Archives of Otolaryngology – Head & Neck Surgery*, 116(4), 424–427. <https://doi.org/10.1001/archotol.1990.01870040046011>
- Jiam, N.T.L., Li, C., & Agrawal, Y. (2016). Hearing loss and falls: A systematic review and meta-analysis. *The Laryngoscope*, 126(11), 2587–2596. <https://doi.org/10.1002/lary.25927>
- Johnson, S.T. (2018). *Falls and fall prevention in community-dwelling older adults*. Digital comprehensive summaries of Uppsala dissertations from the Faculty of Medicine. Uppsala Universitet 1396. Retrieved from <https://www.sakersenior.se/sakersenior/dokument/Fallprevention-Susanna-Tuvemo-Johnson.pdf>
- Kalu, M.E., Vlachantoni, A., & Norman, K.E. (2019). Knowledge about risk factors for falls and practice about fall prevention in older adults amongst physiotherapists in Nigeria. *Physiotherapy Research International*, 24(1), e1742. <https://doi.org/10.1002/pri.1742>
- Kalula, S.Z., Ferreira, M., Swingler, G.H., & Badri, M. (2016). Risk factors for falls in older adults in a South African urban community. *BMC Geriatrics*, 16, 51. <https://doi.org/10.1186/s12877-016-0212-7>
- Kamanji, R. (2016). *Zulu translation and cross-cultural adaptation of the activities-specific balance confidence (ABC) scale*. Research report. University of the Witwatersrand.
- Khow, K.S., & Visvanathan, R. (2017). Falls in the aging population. *Clinics in Geriatric Medicine*, 33(3), 357–368. <https://doi.org/10.1016/j.cger.2017.03.002>
- Lajoie, Y., & Gallagher, S. P. (2004). Predicting falls within the elderly community: Comparison of postural sway, reaction time, the berg balance scale and the activities-specific balance confidence (ABC) scale for comparing fallers and non-fallers. *Archives of Gerontology and Geriatrics*, 38(1), 11–26. [https://doi.org/10.1016/S0167-4943\(03\)00082-7](https://doi.org/10.1016/S0167-4943(03)00082-7)
- Lockhart, T.E., Frames, C.W., Soangra, R., & Lieberman, A. (2019). Effects of obesity and fall risk on gait and posture of community-dwelling older adults. *International Journal of Prognostics and Health Management*, 10(1), 1–10. <https://doi.org/10.36001/ijphm.2019.v10i1.2750>
- Maidan, I., Freedman, T., Tzemah, R., Giladi, N., Mirelman, A., & Hausdorff, J.M. (2014). Introducing a new definition of a near fall: Intra-rater and inter-rater reliability. *Gait & Posture*, 39(1), 645–647. <https://doi.org/10.1016/j.gaitpost.2013.07.123>
- Mak, M.K., Lau, A.L., Law, F.S., Cheung, C.C., & Wong, I.S. (2007). Validation of the Chinese translated activities-specific balance confidence scale. *Archives of Physical Medicine and Rehabilitation*, 88(4), 496–503. <https://doi.org/10.1016/j.apmr.2007.01.018>
- Marques, A.P., Mendes, Y.C., Taddei, U., Pereira, C.A., & Assumpção, A. (2013). Brazilian-Portuguese translation and cross-cultural adaptation of the activities-specific balance confidence (ABC) scale. *Brazilian Journal of Physical Therapy*, 17(2), 170–178. <https://doi.org/10.1590/S1413-35552012005000072>
- McHugh, M.L. (2012). Interrater reliability: The kappa statistic. *Biochemia Medica*, 22(3), 276–282. <https://doi.org/10.11613/BM.2012.031>
- Moiz, J.A., Bansal, V., Noohu, M.M., Gaur, S.N., Hussain, M.E., Anwer, S., & Alghadir, A. (2017). Activities-specific balance confidence scale for predicting future falls in Indian older adults. *Clinical Interventions in Aging*, 12, 645–651. <https://doi.org/10.2147/CIA.S133523>
- Montero-Odasso, M., Van der Velde, N., Martin, F.C., Petrovic, M., Tan, M.P., Ryg, J., Aguilar-Navarro, S., Alexander, N.B., Becker, C., Blain, H., Bourke, R., Cameron, I. D., Camicioli, R., Clemson, L., Close, J., Delbaere, K., Duan, L., Duque, G., Dyer, S.M., Freiburger, E. et al. (2022). World guidelines for falls prevention and management for older adults: A global initiative. *Age and Ageing*, 51(9), afac205. <https://doi.org/10.1093/ageing/afac205>
- Myers, A.M., Fletcher, P.C., Myers, A.H., & Sherk, W. (1998). Discriminative and evaluative properties of the activities-specific balance confidence (ABC) scale. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 53(4), M287–M294. <https://doi.org/10.1093/gerona/53A.4.M287>
- Nagai, K., Yamada, M., Komatsu, M., Tamaki, A., Kanai, M., Miyamoto, T., Tsukagoshi, R., & Tsuboyama, T. (2017). Near falls predict substantial falls in older adults: A prospective cohort study. *Geriatrics & Gerontology International*, 17(10), 1477–1480. <https://doi.org/10.1111/ggi.12898>
- Oh, S.J., Huang, J.J., Hujsak, B., Campo, M., Schwam, Z.G., Cosetti, M.K., Gurley, J., & Kelly, J. (2023). Relevance of the activities-specific balance confidence (ABC) scale to urban community-dwelling adults with diagnosed vestibular dysfunction. *Archives of Physical Medicine and Rehabilitation*, 104(4), 686–689. <https://doi.org/10.1016/j.apmr.2022.09.009>
- Pascoe, M., & Norman, V. (2011). Contextually relevant resources in speech-language therapy and audiology in South Africa – Are there any?. *South African Journal of Communication Disorders*, 58(1), 2–5. <https://doi.org/10.4102/sajcd.v58i1.35>
- Podsiadlo, D., & Richardson, S. (1991). The timed 'up and go': A test of basic functional mobility for frail elderly persons. *Journal of the American Geriatrics Society*, 39(2), 142–148. <https://doi.org/10.1111/j.1532-5415.1991.tb01616.x>
- Powell, L.E., & Myers, A.M. (1995). The activities-specific balance confidence (ABC) scale. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 50A(1), M28–M34. <https://doi.org/10.1093/gerona/50A.1.M28>
- Reelink, M.F., Van Iersel, M.B., Kessels, R.P.C., & Olde Rikkert, M.G.M. (2009). The influence of fear of falling on gait and balance in older people. *Age and Ageing*, 38(4), 435–440. <https://doi.org/10.1093/ageing/afp066>
- Rogers, C. (2021). Audiologists should not fail with falls: A call to commit to prevention of falls in older adults. *South African Journal of Communication Disorders*, 68(1), a841. <https://doi.org/10.4102/sajcd.v68i1.841>
- Sahu, U.R., & Kulkarni, V.N. (2017). Translation and cross-cultural adaptation of the activities-specific balance confidence (ABC) scale for use in Maharashtra. *Journal of The Indian Academy of Geriatrics*, 13(2), 73–76. <https://doi.org/10.35262/jiag.v13i2.73-76>
- Salari, N., Darvishi, N., Ahmadipannah, M., Shohaimi, S., & Mohammadi, M. (2022). Global prevalence of falls in the older adults: A comprehensive systematic review and meta-analysis. *Journal of Orthopaedic Surgery and Research*, 17, 334. <https://doi.org/10.1186/s13018-022-03222-1>
- Sakakibara, B.M., Miller, W.C., & Backman, C.L. (2011). Rasch analyses of the activities-specific balance confidence scale with individuals 50 years and older with lower-limb amputations. *Archives of Physical Medicine and Rehabilitation*, 92(8), 1257–1263. <https://doi.org/10.1016/j.apmr.2011.03.013>
- Salbach, N.M., Mayo, N.E., Hanley, J.A., Richards, C.L., & Wood-Dauphinee, S. (2006). Psychometric evaluation of the original and Canadian French version of the activities-specific balance confidence scale amongst people with stroke. *Archives of Physical Medicine and Rehabilitation*, 87(12), 1597–1604. <https://doi.org/10.1016/j.apmr.2006.08.336>
- Schepens, S., Goldberg, A., & Wallace, M. (2010). The short version of the activities-specific balance confidence (ABC) scale: Its validity, reliability, and relationship to balance impairment and falls in older adults. *Archives of Gerontology and Geriatrics*, 51(1), 9–12. <https://doi.org/10.1016/j.archger.2009.06.003>
- Schott, N. (2008). German adaptation of the 'Activities-specific Balance Confidence (ABC) scale' for the assessment of falls-related self-efficacy. *Zeitschrift für Gerontologie und Geriatrie*, 41(6), 475–485. <https://doi.org/10.1007/s00391-007-0504-9>
- Seamon, B.A., Kautz, S.A., & Velozo, C.A. (2019). Rasch analysis of the Activities-Specific Balance Confidence scale in individuals poststroke. *Archives of Rehabilitation Research and Clinical Translation*, 1(3–4), 100028. <https://doi.org/10.1016/j.arct.2019.100028>
- Solanki, G., Kelly, G., Cornell, J., Daviaud, E., & Geffen, L. (2019). Population ageing in South Africa: Trends, impact, and challenges for the health sector. *South African Health Review*, 1, 175–182. Retrieved from <https://hdl.handle.net/10520/EJC-1d2b0b4f5a>
- Statistics South Africa. (2011). *Census 2011: Census in brief*. Retrieved from [https://www.statssa.gov.za/census/census\\_2011/census\\_products/Census\\_2011\\_Census\\_in\\_brief.pdf](https://www.statssa.gov.za/census/census_2011/census_products/Census_2011_Census_in_brief.pdf)
- Statistics South Africa. (2018). *General household survey: 2018. Statistical release P0318*. Retrieved from <http://www.statssa.gov.za/publications/P0318/P03182018.pdf>
- Statistics South Africa. (2020a). *Mid-year population estimates. Statistical release P0302*. Retrieved from <https://www.statssa.gov.za/publications/P0302/P03022020.pdf>



- Statistics South Africa. (2020b). *National households travel survey 2020*. Statistical release P0320. Retrieved from <http://www.statssa.gov.za/publications/P0320/P03202020.pdf>
- Tavakol, M., & Dennick, R. (2011). Making sense of Cronbach's alpha. *International Journal of Medical Education*, 2, 53–55. <https://doi.org/10.5116/ijme.4dfb.8dfd>
- United Nations. (2015a). *World population ageing 2015 – Highlights*. Retrieved from [https://www.un.org/en/development/desa/population/publications/pdf/ageing/WPA2015\\_Highlights.pdf](https://www.un.org/en/development/desa/population/publications/pdf/ageing/WPA2015_Highlights.pdf)
- United Nations. (2015b). *World population prospects: The 2015 revision*. Retrieved from [https://population.un.org/wpp/Publications/Files/Key\\_Findings\\_WPP\\_2015.pdf](https://population.un.org/wpp/Publications/Files/Key_Findings_WPP_2015.pdf)
- United Nations. (2019). *World population ageing 2019 – Highlights*. Retrieved from <https://www.un.org/en/development/desa/population/publications/pdf/ageing/WorldPopulationAgeing2019-Highlights.pdf>
- Van den Berg, V.L. (2016). Still lost in translation: Language barriers in South African health care remain. *South African Family Practice*, 58(6), 229–231. <https://doi.org/10.1080/20786190.2016.1223795>
- Van Heuvelen, M.J., Hochstenbach, J., De Greef, M.H., Brouwer, W.H., Mulder, T., & Scherder, E. (2005). Is the activities-specific balance confidence scale suitable for Dutch older persons living in the community?. *Tijdschrift voor Gerontologie en Geriatrie*, 36(4), 146–154. <https://doi.org/10.1007/BF03074728>
- Van Rie, K.J., Kanji, A., & Naude, A. (2022). Professional guidelines and reported practice of audiologists performing fall risk assessment with older adults: A systematic review. *American Journal of Audiology*, 31(1), 243–260. [https://doi.org/10.1044/2021\\_AJA-21-00148](https://doi.org/10.1044/2021_AJA-21-00148)
- World Health Organization. (2001). *International classification of functioning, disability and health*. Retrieved from <https://apps.who.int/iris/bitstream/handle/10665/42407/9241545429-eng.pdf?sequence=1&isAllowed=y>
- World Health Organization. (2002). *Active ageing: A policy framework*. Retrieved from <https://extranet.who.int/agefriendlyworld/wp-content/uploads/2014/06/WHO-Active-Ageing-Framework.pdf>
- World Health Organization. (2016). *Process of translation and adaptation of instruments*. Retrieved from <https://www.mhinnovation.net/sites/default/files/files/WHO%20Guidelines%20on%20Translation%20and%20Adaptation%20of%20Instruments.docx>
- World Health Organization. (2021). *Falls*. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/falls>
- World Health Organization. (2022). *Ageing and health*. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>
- World Medical Association. (2013). *WMA declaration of Helsinki: Ethical principles for medical research involving human subjects*. Retrieved from <https://www.wma.net/policies-post/wma-declaration-of-helsinki-ethical-principles-for-medical-research-involving-human-subjects/>

Appendices starts on the next page →

## Appendix 1: Case history questionnaire.

### Participant particulars

Participant number: \_\_\_\_\_ Gender: \_\_\_\_\_ Age: \_\_\_\_\_

Home language: \_\_\_\_\_ Date: \_\_\_\_\_

Please complete the following (*please check [v] the box that applies to you*)

#### Marital status

Single  Married  Divorced  Widowed

#### Highest level of education

No formal education  Grade 1–7  Grade 8–9  
 Grade 10–12  College  University

### Balance history

- Do you have a fear of falling?
  - Yes
  - No
- Do you use walking aids indoors? If yes, please describe.
  - Yes \_\_\_\_\_
  - No
- Do you use walking aids outdoors? If yes, please describe.
  - Yes \_\_\_\_\_
  - No
- Do you need physical assistance when walking outside? If yes, please explain what assistance you require, for example: holding onto someone's arm.
  - Yes \_\_\_\_\_
  - No
- Have you felt unsteady on your feet in the last 6 months?
  - Yes
  - No
- Have you fallen anytime in the last 6 months? If yes, please indicate how many times you have fallen.
  - Yes \_\_\_\_\_
  - No
- How many times do you exercise per week?
  - Less than three times
  - Three times or more
- Please list what exercise you do below and for how long.
  - \_\_\_\_\_
  - \_\_\_\_\_
  - \_\_\_\_\_

### Medical history

- Please list any allergies that you have.
  - \_\_\_\_\_
  - \_\_\_\_\_
  - \_\_\_\_\_
- Please list any medical conditions that you have.
  - \_\_\_\_\_
  - \_\_\_\_\_

- \_\_\_\_\_
- Please list the medication that you are currently taking.
    - \_\_\_\_\_
    - \_\_\_\_\_
    - \_\_\_\_\_
  - Please list any surgeries or operations that you have had.
    - \_\_\_\_\_
    - \_\_\_\_\_
    - \_\_\_\_\_
  - Have you had any previous medical examination or testing done? (Hearing, vision, x-rays, head scans etc.). If yes, please provide details as well as a summary of the results if possible.
    - Yes \_\_\_\_\_
    - No

### Family history

- Please indicate if you have a family history of any of the following by ticking the box.
  - Hearing loss
  - Dizziness
  - High blood pressure
  - Low blood pressure
  - Headaches
  - Migraines
  - Diabetes
  - Panic attacks

## Appendix 2: Sepedi version of the ABC (ABC-S) scale.

Kelo ya mešongwana- yeo e ikgethilego ya boitshepo bja palantshe (ABC)

Go mošongwana o mongwe le o mongwe woo o latelago, kgopela o laetše maemo a gago a boitshepo ka go kgetha nomoro ya maleba go ya le ka maemo a kelo:

0% 10 20 30 40 50 60 70 80 90 100%

*go hloka boithsepo*

*go ba le boitshepo*

'Ona le boitshepo bjo bo kakang ba gore o ka se lahlegele ke palantshe goba go tšhekinyega ge o...

- ...ge o sepela ka gae? \_\_\_\_\_%
- ...o eya godimo le fase disetepiseng? \_\_\_\_\_%
- ...ge o khunama o topa seetabolaong fase ge o tloga ka pele ga bo bolokelo bja phaahlo? \_\_\_\_\_%
- ...ge o obeletša kotikoti ye nnyane godimo ga šelefe yeo o kgonago go e bona? \_\_\_\_\_%
- ...go tsatsampele gore o kgone go fihlelela seo se lego ka godimo ga hlogo ya gago? \_\_\_\_\_%
- ...go namela setulo go ka kgona go fihlelela se sengwe? \_\_\_\_\_%
- ...go swiela? \_\_\_\_\_%
- ...go sepela ka ntle o eya ka jareteng mo go emago sefatanaga? \_\_\_\_\_%
- ...go namela le go fologa sefatanaga? \_\_\_\_\_%

10. ...o tloga bo dulong bja difatanaga go ya mmolong/mabenkeleng? \_\_\_\_%
11. ...go namela le go fologa mototolo? \_\_\_\_%
12. ...o sepela ka mmolong/mabenkeleng moo batho e lego ba bantši ba fetafeta kgauswi le wena? \_\_\_\_%
13. ...go hlakhlakana le batho ge o sepela ka mmolong/mabenkeleng? \_\_\_\_%
14. ... o namela setepisitshepetšo o itshwareleditše? \_\_\_\_%
15. ... o namela goba o fologa setepisitshepetšo o swere merwalo yeo e ka go paledišago go itshwareletša? \_\_\_\_%

16. ...ge o sepela ka ntle go dimo ga phoka? \_\_\_\_%

### Instructions for scoring

The ABC is an 11-point scale and ratings should consist of whole numbers (0–100) for each item. Total the ratings (possible range = 0–1600) and divide by 16 to get each person's ABC score.

Total score: \_\_\_\_\_